www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

Biotin Human ACE-2 Protein (C-His-Avi)

Catalog Number:	601103, 601104
Size:	25 ug, 100 ug
Target Name:	ACE2, angiotensin converting Enzyme 2, ACEH
Regulatory Status:	RUO

PRODUCT DETAILS


Application:	ELISA, BLI
Format:	Liquid, Biotinylated
Expression Host:	CHO
Species:	Human
Accession Number:	Q9BYF1
Sources:	Human ACE2 protein (Gln18-Ser740) with C-terminus His-Avi tag is expressed in CHO cells. This protein was site-specifically labeled with Biotin by BirA ligase.
Molecular Weight:	This protein has a predicted molecular weight of 87.4 kDa. Under DTT-reducing conditions, the protein migrates at approximately 90-100 kDa on SDS-PAGE.
Affinity Tag:	C-His-Avi
Purity:	>95% based on SDS-PAGE under reducing condition
Formulation:	1xPBS buffer, pH7.4, 0.22 μm filtered
Endotoxin level:	Not tested
Protein Concentration:	25μg size is bottled at 0.2mg/mL concentration. 100 μg size is supplied at a lot-specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 μm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ –70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.
Recommended Usage:	For detection, use a secondary reagent with this product.

BACKGROUND INFORMATION

Angiotensin-converting enzyme 2 (ACE2) is a type I transmembrane zinc metalloprotease with about 60% homology to ACE, consisting of 805 amino acids including a signal peptide, catalytic domain, membrane anchor, and cytoplasmic tail. ACE2 regulates the renin-angiotensin system by converting angiotensin I to angiotensin 1-9 and angiotensin II to the vasodilator angiotensin 1-7, thus counteracting vasoconstriction and playing a key role in cardiovascular and renal function. ACE2 also serves as the functional receptor for human coronaviruses SARS-CoV, SARS-CoV-2, and HCoV-NL63, facilitating viral entry. Its expression is high in lung,

heart, kidney, intestine, testis, and vascular cells. Beyond cardiovascular regulation, ACE2 is involved in respiratory disease pathogenesis, including acute respiratory distress syndrome, and has been linked to diabetes and hypertension. Additionally, its homolog collectrin is implicated in amino acid transport and genetic disorders like Hartnup disease.

PRODUCT DATA



SARS-CoV-2 Spike RBD Protein (C-Fc, Catalog # 602501) is coated at 2 μg/mL (200 ng/well). Biotinylated human ACE2 with C-Avi-His can bind RBD in the dose dependent manner. The EC50 is in the range of 5-20 ng/mL

This product is supplied subject to the terms and conditions at www.innocyto.com.cn/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.