www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

iF647 Anti-Human CD137 (4-1BB) Antibody

Catalog Number:	108003, 108004
Size:	25 tests, 100 tests
Target Name:	CD137, 4-1BB, ILA, TNFRSF9
Regulatory Status:	RUO


PRODUCT DETAILS

Clone:	4B4-1
Application:	Flow Cytometry
Reactivity:	Human
Format:	iF647
Isotype:	Mouse IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. iF647 has an excitation max at 656 nm and an emission max at 670 nm.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	301413
RRID:	AB_3738814

BACKGROUND INFORMATION

CD137 is a 39 kDa type I transmembrane protein and a member of the Tumor Necrosis Factor Receptor (TNFR) superfamily. It is expressed on activated T cells, NKT cells, and NK cells, where it functions as a co-stimulatory molecule that promotes T cell proliferation, survival, and monocyte activation through its interaction with 4-1BB ligand. CD137 plays a key role in anti-tumor immunity. Agonistic anti-CD137 antibodies enhance tumor-specific $CD8^+$ T cell responses, increase NK and dendritic cell function, and improve antibody-dependent cellular cytotoxicity (ADCC) when used with tumor-targeting monoclonal antibodies. Its expression can be upregulated on NK cells after CD16 ligation, although its effect on NK cell cytotoxicity can vary between human and murine systems. Overall, CD137 is an important immunomodulatory target in cancer immunotherapy.

PRODUCT DATA



Anti-human CD3 and Anti-Human CD28 stimulated human peripheral blood lymphocytes was stained with iF647 Anti-Human CD137 clone 4B4-1 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com.cn/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.