www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

APC Anti-mouse CD19 Antibody

Catalog Number:	201909, 201910
Size:	25 μg, 100 μg
Target Name:	CD19, B4, CVID3, Leu-12, MGC12802
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	1D3
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	APC
Isotype:	Rat IgG2a
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.3 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm.
Excitation Laser:	Red Laser (633 nm)

BACKGROUND INFORMATION

CD19, also known as B-lymphocyte antigen CD19, B4, Leu-12, or CVID3, is a type I transmembrane glycoprotein encoded by the CD19 gene. It is expressed throughout B cell development from the pro-B stage to mature B cells and is present on most B lineage cells. CD19 functions as an adaptor protein, recruiting cytoplasmic signaling molecules to the membrane, and forms part of the CD19/CD21/CD81 complex, which lowers the activation threshold for B cell receptor (BCR) signaling. Because of its consistent expression on B cells, CD19 serves as a key biomarker for B cell development and is widely used in the diagnosis of B cell lymphomas and leukemias. It is also a major therapeutic target in immunotherapies, including CAR T-cell therapy for B cell malignancies.