www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

FITC Anti-Mouse/Human CD44 Antibody

Catalog Number:	202107, 202108
Size:	50 μg, 500 μg
Target Name:	CD44, Epican, Extracellular matrix receptor III (ECMR-III), Phagocytic glycoprotein 1 (PGP-1)
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	IM7
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	FITC
Isotype:	Rat IgG2b
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.3 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. FITC has an excitation max at 495 nm and an emission max at 519 nm.
Excitation Laser:	Blue Laser (488 nm)
RRID:	AB_3739079

BACKGROUND INFORMATION

CD44 is an 80–100 kDa type I transmembrane glycoprotein and a member of the cell adhesion molecule family. It acts as a receptor for hyaluronic acid (HA) and mediates cell-cell interactions, adhesion, migration, and signal transduction. CD44 exists in multiple isoforms generated through alternative splicing, which contributes to its functional diversity.