www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

iF488 Anti-human CD282 (TLR2) Antibody

Catalog Number:	114505, 114506
Size:	25 tests, 100 tests
Target Name:	CD282, TLR2, Toll like receptor 2, TIL4
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	282AM2a
Application:	Flow Cytometry
Reactivity:	Human
Format:	iF488
Isotype:	Mouse IgG2a
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. iF488 has an excitation max at 491 nm and an emission max at 516 nm.
Excitation Laser:	Blue Laser (488 nm)
Isotype Controls:	301505
Antibody Family:	Human Antibodies

BACKGROUND INFORMATION

Toll-like receptor 2 (TLR2) is a pattern recognition receptor expressed on innate immune cells such as monocytes, macrophages, dendritic cells, and some epithelial cells. It plays a key role in host defense by recognizing conserved microbial components and initiating inflammatory signaling pathways. Upon activation, TLR2 triggers downstream signaling through adaptor proteins like MyD88, leading to NF-κB activation and production of pro-inflammatory cytokines.


Structurally, TLR2 is a type I transmembrane protein composed of an extracellular domain rich in leucine-rich repeats (LRRs) that mediate ligand recognition, a single transmembrane helix, and an intracellular Toll/IL-1 receptor (TIR) domain responsible for signal transduction. TLR2 typically forms heterodimers with TLR1 or TLR6, which expands its ligand specificity.

TLR2 recognizes a broad range of ligands, including bacterial lipoproteins, peptidoglycan, lipoteichoic acid from Gram-positive

bacteria, and certain fungal and viral components. Endogenous ligands released during tissue damage can also activate TLR2, linking it to sterile inflammation.

In disease, dysregulated TLR2 signaling contributes to chronic inflammatory conditions, autoimmune diseases, and sepsis. It also plays roles in cancer by influencing tumor-associated inflammation. Therapeutically, TLR2 is being explored as both a target and a tool, with agonists used as vaccine adjuvants and antagonists investigated to reduce harmful inflammation.

PRODUCT DATA



Human peripheralmonocyteswerestainedwithiF488 anti-Human CD282 (TLR2) clone 282AM2a (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com.cn/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.