www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

PB Anti-Mouse/Human KLRG1 Antibody

Catalog Number:	204217, 204218
Size:	25 μg, 100 μg
Target Name:	KLRG1, MAFA, 2F1-Ag
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	2F1
Application:	Flow Cytometry
Reactivity:	Mouse, Human
Format:	PB
Isotype:	Syrian Hamster IgG
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	0.2 mg/mL
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.3 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. Pacific Blue has an excitation max at 404 nm and an emission max at 455 nm.
Excitation Laser:	Violet laser (405nm)
Isotype Control:	303912

BACKGROUND INFORMATION

KLRG1 (Killer Cell Lectin-Like Receptor Subfamily G Member 1) is an inhibitory receptor expressed on several immune cell populations, particularly natural killer (NK) cells and subsets of T lymphocytes. It is commonly used as a marker of terminal differentiation or functional maturation in effector CD8+ T cells and NK cells. KLRG1 plays an important role in regulating immune responses by delivering inhibitory signals that limit excessive activation, thereby helping maintain immune homeostasis during infections and inflammatory responses.

Structurally, KLRG1 is a type II transmembrane glycoprotein belonging to the C-type lectin-like receptor family. The protein contains a short N-terminal cytoplasmic domain, a single transmembrane region, and an extracellular C-type lectin-like domain responsible for ligand binding. Unlike classical C-type lectins, KLRG1 does not bind carbohydrates but instead recognizes protein ligands. The cytoplasmic tail contains an immunoreceptor tyrosine-based inhibitory motif (ITIM). When the receptor engages its ligand, this ITIM becomes phosphorylated and recruits phosphatases such as SHP-1 or SHP-2, which suppress downstream signaling pathways


involved in cellular activation.

The primary ligands for KLRG1 are members of the classical cadherin family, including E-cadherin, N-cadherin, and R-cadherin. These molecules are widely expressed on epithelial cells and other tissue types. Interaction between KLRG1 on immune cells and cadherins on target cells transmits inhibitory signals that dampen cytotoxic activity and cytokine production. This interaction is thought to help prevent excessive immune-mediated tissue damage, particularly at epithelial surfaces where cadherins are abundant.

KLRG1 has been studied extensively in the context of infections, aging, and cancer. During viral or bacterial infections, a population of KLRG1high CD8+ T cells often emerges; these cells represent short-lived effector cells that exhibit strong cytotoxic activity but limited proliferative capacity. In chronic infections and aging, increased frequencies of KLRG1-expressing T cells are associated with immune senescence and reduced immune responsiveness. In cancer, KLRG1-mediated inhibitory signaling may contribute to reduced anti-tumor activity of NK and T cells within the tumor microenvironment.

Because of its role as an immune inhibitory receptor, KLRG1 has attracted interest as a potential target for immunotherapy. Blocking KLRG1 interactions with cadherins could enhance the cytotoxic function of NK cells and CD8+ T cells, thereby improving anti-tumor immune responses. Conversely, enhancing KLRG1 signaling might be beneficial in conditions where immune responses need to be restrained, such as autoimmune or inflammatory diseases. In addition, KLRG1 expression is widely used as a biomarker in immunological research to identify differentiated effector T cells and to track immune responses in infection and vaccination models.

PRODUCT DATA



C57BL/6 Mouse splenocytes were stained with iF647 Anti-Mouse NK1.1 clone PK136 and PB Anti- Mouse/Human KLRG1 clone 2F1 (left) or iF647 Anti-Mouse NK1.1 clone PK136 only (right).

This product is supplied subject to the terms and conditions at www.innocyto.com.cn/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.