www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

APC Anti-Human/Mouse/Rat CD278 (ICOS) Antibody

Catalog Number:	111209, 111210
Size:	25 ug, 100 ug
Target Name:	CD278, ICOS, Inducible Costimulatory molecule
Regulatory Status:	RUO


PRODUCT DETAILS

Clone:	C398.4A
Application:	Flow Cytometry
Reactivity:	Human, mouse, rat
Format:	APC
Isotype:	Armenian Hamster IgG
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	300514
Recommended usage:	For flow cytometric staining, it is recommended to use less than 0.5 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm.

BACKGROUND INFORMATION

ICOS is a 47–57 kDa costimulatory molecule homologous to CD28/CTLA-4 and is expressed on activated T cells and a subset of thymocytes. Engagement of ICOS enhances T cell proliferation, promotes cytokine production such as IL-4 and IL-10, and plays an important role in humoral immunity and Th2 cell differentiation.

PRODUCT DATA



Human peripheral blood lymphocytes stimulated with Anti-Human CD3/CD28for3-dayswerestained with APC Anti-Human/Mouse/Rat CD278 (ICOS) clone C398.4A (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com.cn/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.