www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

PE/Cyanine5.5 Anti-mouse CD279 (PD-1) Antibody

Catalog Number:	205021, 205022
Size:	25 µg, 100 µg
Target Name:	CD279, Programmed Death-1, PD 1, PDCD1, PD-1
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	29F.1A12
Application:	Flow Cytometry
Reactivity:	Mouse
Format:	PE/Cyanine5.5
Isotype:	Rat IgG2a
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	0.2 mg/mL
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use less than 0.25 µg of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. PE/Cyanine5.5 has an excitation max at 565 nm and an emission max at 670 nm.
Excitation Laser:	Blue Laser (488 nm) Green/Yellow laser (532/561nm)
Isotype Control:	303515

BACKGROUND INFORMATION

Mouse CD279, more commonly known as programmed cell death protein 1 (PD-1), is an inhibitory immune checkpoint receptor expressed primarily on activated T cells, as well as B cells and some myeloid populations. It plays a critical role in maintaining peripheral tolerance and preventing excessive immune activation by downregulating T cell responses during chronic antigen exposure, such as infection or inflammation. PD-1 is rapidly induced following T cell receptor engagement and acts as a key regulator of immune homeostasis.

Structurally, PD-1 is a type I transmembrane protein belonging to the immunoglobulin superfamily. It contains a single extracellular IgV-like domain responsible for ligand binding, a transmembrane region, and a cytoplasmic tail with immunoreceptor tyrosine-based inhibitory (ITIM) and switch (ITSM) motifs. Upon ligand engagement, these motifs recruit phosphatases such as SHP-2, which attenuate proximal T cell receptor signaling pathways.


The primary ligands of PD-1 are PD-L1 (CD274) and PD-L2 (CD273), which are expressed on antigen-presenting cells and various

non-hematopoietic tissues. Interaction of PD-1 with its ligands suppresses T cell proliferation, cytokine production, and survival, promoting an exhausted T cell phenotype during chronic immune stimulation.

PD-1 signaling is implicated in chronic infections, cancer, and autoimmune diseases. In tumors, PD-1-mediated inhibition allows cancer cells to evade immune surveillance by suppressing anti-tumor T cell activity.

Therapeutically, blockade of the PD-1/PD-L1 axis using monoclonal antibodies has revolutionized cancer immunotherapy by restoring T cell function. Conversely, enhancing PD-1 signaling may be beneficial in treating autoimmune diseases and preventing transplant rejection, making it a versatile target in immune modulation.

PRODUCT DATA



C57BL/6 Mouse splenocytes stimulated with Anti-Mouse CD3/CD28 for 2-days were stained with PE/Cyanine5.5 Anti-Mouse CD279 (PD-1) clone 29F.1A12 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com.cn/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.