www.innocyto.com.cn

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

In Vivo Star Rat IgG2a Isotype Control Antibody

Catalog Number:	570801, 570802, 570803
Size:	1 mg, 5 mg, 25 mg
Target Name:	Rat IgG2a Isotype Control Antibody
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	IVSR2a
Application:	An isotype-matched negative control for Rat IgG2a, kappa antibodies used in ELISA, Western Blot (WB), Flow Cytometry (Flow), Immunoprecipitation (IP), Immunohistochemistry (Paraffin) (IHC (P)), Immunohistochemistry (Frozen) (IHC (F)), and in vivo animal model research.
Reactivity:	N/A
Format:	Liquid
Isotype:	Rat IgG2a, κ
Antibody Type:	Recombinant
Formulation:	0.2 μM filtered solution of 1x PBS.
Endotoxin Level:	Less than 1 EU/mg of protein as determined by LAL method.

BACKGROUND INFORMATION

Rat IgG isotype control antibodies are non-specific immunoglobulins used as negative controls in immunological and cell-based assays. They are matched to the species, immunoglobulin class or subclass, and, when applicable, light-chain type of the primary antibody being tested. Rat IgG antibodies can be classified into several isotypes based on their heavy-chain composition, including IgG1, IgG2a, IgG2b, and IgG2c, depending on the rat strain and nomenclature used. Their light chains are primarily kappa (κ) or lambda (λ).

Rat IgG antibodies are glycoproteins with a typical molecular weight of approximately 150 kDa and consist of two identical heavy chains and two identical light chains. The four chains are connected by disulfide bonds, forming the characteristic Y-shaped antibody structure. The two antigen-binding sites are located within the Fab regions at the ends of the antibody arms, while the Fc region is responsible for interactions with Fc receptors and other immune components. For an isotype control, the antibody is selected to have the same or closely matched isotype and structural characteristics as the test antibody but lacks specific binding to the target antigen.

Rat IgG isotype controls are widely used in flow cytometry, immunohistochemistry, immunofluorescence, and other antibody-based assays to help evaluate nonspecific antibody binding and background staining. An appropriate isotype control should ideally match the primary antibody in species, isotype, conjugate, concentration, and antibody-to-protein ratio. By providing a measure of nonspecific interactions, rat IgG isotype controls can help researchers distinguish true antigen-specific signals from background caused by Fc-receptor interactions, cellular autofluorescence, or nonspecific antibody binding.